

Wavelets for reinforced function spaces on cellular domains

Benjamin Scharf

Friedrich Schiller University Jena

June 20, 2012

Table of contents

- 1 Wavelets bases in function spaces on domains - the classical cases
 - Known results - wavelets on domains
 - Function spaces on domains
 - Traces on the boundary of cubes
- 2 Wavelets bases in reinforced function spaces - the exceptional values
 - The situation for cubes Q in the exceptional cases
 - Wavelet bases for reinforced function spaces on cubes Q
 - An example - $W_2^{1,\text{rinf}}(Q)$

Wavelets for cubes (i)

Let Q be the unit cube in \mathbb{R}^n .

Theorem (Triebel 2008 - Theorem 6.30 for spaces $F_{p,q}^s(Q)$)

Let

$$1 < p < \infty, u, k \in \mathbb{N}_0, u > k \text{ and } k - \frac{m}{p} \notin \mathbb{N}_0$$

for $m = 1, \dots, n$.

Then there is an oscillating u -wavelet system Φ which is a Riesz basis in the Sobolev space $W_p^k(Q)$,

Wavelets for cubes (i)

Let Q be the unit cube in \mathbb{R}^n .

Theorem (Triebel 2008 - Theorem 6.30 for spaces $F_{p,q}^s(Q)$)

Let $1 < p < \infty, u, k \in \mathbb{N}_0, u > k$ and $k - \frac{m}{p} \notin \mathbb{N}_0$

for $m = 1, \dots, n$.

Then there is an oscillating u -wavelet system Φ which is a Riesz basis in the Sobolev space $W_p^k(Q)$, i.e.: An element $f \in D'(\Omega)$ belongs to $W_p^k(Q)$ if, and only if, it can be represented as

$$f = \sum_{j=0}^{\infty} \sum_{r=1}^{N_j} \lambda_r^j(f) 2^{-\frac{jn}{2}} \Phi_r^j \quad (1)$$

with λ in the sequence space $w_p^k(Q)$ and the convergence is unconditional in $W_p^k(Q)$. The representation (1) is unique with

$$\lambda_r^j(f) = 2^{jn/2} (f, \Phi_r^j).$$

The spaces $W_p^k(\mathbb{R}^n)$ and $W_p^k(\Omega)$

Definition

Let $1 < p < \infty$ and $k \in \mathbb{N}_0$. Then

$$W_p^k(\mathbb{R}^n) := \{f \in D'(\mathbb{R}^n) : D^\alpha f \in L_p(\mathbb{R}^n) \text{ for } |\alpha| \leq k\}$$

with

$$\|f|W_p^k(\mathbb{R}^n)\| = \sum_{|\alpha| \leq k} \|D^\alpha f|L_p(\mathbb{R}^n)\|.$$

Let Ω be an arbitrary domain, i. e. open set in \mathbb{R}^n . Then

$$W_p^k(\Omega) := \{f \in D'(\Omega) : f = g|_\Omega \text{ for some } g \in W_p^k(\mathbb{R}^n)\},$$

$$\|f|W_p^k(\Omega)\| = \inf \|g|W_p^k(\mathbb{R}^n)\|$$

where the infimum is taken over all $g \in W_p^k(\mathbb{R}^n)$ with $g|_\Omega = f$.

The spaces $W_p^k(\mathbb{R}^n)$ and $W_p^k(\Omega)$

Definition

Let $1 < p < \infty$ and $k \in \mathbb{N}_0$. Then

$$W_p^k(\mathbb{R}^n) := \{f \in D'(\mathbb{R}^n) : D^\alpha f \in L_p(\mathbb{R}^n) \text{ for } |\alpha| \leq k\}$$

with

$$\|f|W_p^k(\mathbb{R}^n)\| = \sum_{|\alpha| \leq k} \|D^\alpha f|L_p(\mathbb{R}^n)\|.$$

Let Ω be an arbitrary domain, i. e. open set in \mathbb{R}^n . Then

$$W_p^k(\Omega) := \{f \in D'(\Omega) : f = g|_\Omega \text{ for some } g \in W_p^k(\mathbb{R}^n)\},$$

$$\|f|W_p^k(\Omega)\| = \inf \|g|W_p^k(\mathbb{R}^n)\|$$

where the infimum is taken over all $g \in W_p^k(\mathbb{R}^n)$ with $g|_\Omega = f$.

An equivalent characterization for “good” Ω is given by

$$f \in W_p^k(\Omega) \Leftrightarrow \sum_{|\alpha| \leq k} \|D^\alpha f|L_p(\Omega)\| < \infty \text{ (equivalent norms)}$$

The spaces $w_p^k(\mathbb{Z}^\Omega)$

Let

$$\mathbb{Z}^\Omega := \left\{ x_\ell^j \in \Omega : j \in \mathbb{N}_0, \ell = 1, \dots, N_j \right\},$$

(if Ω bounded, typically $N_j \sim 2^{jn}$), such that for some $c_1 > 0$

$$|x_\ell^j - x_{\ell'}^j| \geq c_1 2^{-j}, \quad j \in \mathbb{N}_0, \ell \neq \ell'.$$

The spaces $w_p^k(\mathbb{Z}^\Omega)$

Let

$$\mathbb{Z}^\Omega := \left\{ x_\ell^j \in \Omega : j \in \mathbb{N}_0, \ell = 1, \dots, N_j \right\},$$

(if Ω bounded, typically $N_j \sim 2^{jn}$), such that for some $c_1 > 0$

$$|x_\ell^j - x_{\ell'}^j| \geq c_1 2^{-j}, \quad j \in \mathbb{N}_0, \ell \neq \ell'.$$

We can introduce the sequence spaces $w_p^k(\mathbb{Z}^\Omega)$ adapted to \mathbb{Z}^Ω : Let

$$\|\lambda\|_{w_p^k(\mathbb{Z}^\Omega)} := \left\| \left(\sum_{j=0}^{\infty} \sum_{\ell=1}^{N_j} 2^{2jsq} |\lambda_\ell^j \cdot \chi_{j,\ell}(\cdot)|^2 \right)^{\frac{1}{2}} \right\|_{L_p(\Omega)},$$

where $\chi_{j,\ell}$ is the characteristic function of $B(x_\ell^j, c2^{-j})$ resp. $\bar{\Omega} \cap B(x_\ell^j, c2^{-j})$ with an arbitrary constant $c > 0$.

Wavelets for function spaces on domains Ω (i)

Let $u \in \mathbb{N}_0$.

Then

$$\Phi = \left\{ \Phi_\ell^j : j \in \mathbb{N}_0, \ell = 1, \dots, N_j \right\} \subset C^u(\Omega)$$

is called a u -wavelet system in $\bar{\Omega}$ (adapted to \mathbb{Z}^Ω) if it fulfils

- support conditions: For some $c_3 > 0$ it holds

$$\text{supp } \Phi_\ell^j \subset B(x_\ell^j, c_3 2^{-j}) \cap \bar{\Omega}, \quad j \in \mathbb{N}_0, \ell = 1, \dots, N_j,$$

Wavelets for function spaces on domains Ω (i)

Let $u \in \mathbb{N}_0$.

Then

$$\Phi = \left\{ \Phi_\ell^j : j \in \mathbb{N}_0, \ell = 1, \dots, N_j \right\} \subset C^u(\Omega)$$

is called a u -wavelet system in $\bar{\Omega}$ (adapted to \mathbb{Z}^Ω) if it fulfils

- support conditions: For some $c_3 > 0$ it holds

$$\text{supp } \Phi_\ell^j \subset B(x_\ell^j, c_3 2^{-j}) \cap \bar{\Omega}, j \in \mathbb{N}_0, \ell = 1, \dots, N_j,$$

- derivative conditions: For some $c_4 > 0$ and all $\alpha \in \mathbb{N}_0^n$ with $0 \leq |\alpha| \leq u$ we have

$$\left| D^\alpha \Phi_\ell^j(x) \right| \leq c_4 2^{j \frac{n}{2} + j|\alpha|}, x \in \Omega, j \in \mathbb{N}_0, \ell = 1, \dots, N_j.$$

Wavelets for function spaces on domains Ω (ii)

Additionally, the u-wavelet system is called oscillating if it fulfils

- (substitute) moment conditions: Let c_5 and $c_6 < c_7$ be constants such that

$$\text{dist}(B(x_\ell^0, c_3), \partial\Omega) \geq c_6, \text{ for } \ell = 1, \dots, N_0 \text{ and}$$

$$\left| \int_{\Omega} \psi(x) \Phi_\ell^j(x) dx \right| \leq c_5 2^{-j\frac{n}{2}-ju} \|\psi\|_{C^u(\Omega)} \text{ for all } \psi \in C^u(\Omega)$$

for all j and ℓ with $\text{dist}(B(x_\ell^j, c_3), \partial\Omega) \notin (c_6 2^{-j}, c_7 2^{-j})$.

Wavelets for function spaces on domains Ω (ii)

Additionally, the u-wavelet system is called oscillating if it fulfils

- (substitute) moment conditions: Let c_5 and $c_6 < c_7$ be constants such that

$$\text{dist}(B(x_\ell^0, c_3), \partial\Omega) \geq c_6, \text{ for } \ell = 1, \dots, N_0 \text{ and}$$

$$\left| \int_{\Omega} \psi(x) \Phi_\ell^j(x) dx \right| \leq c_5 2^{-j \frac{n}{2} - ju} \|\psi\| C^u(\Omega) \text{ for all } \psi \in C^u(\Omega)$$

for all j and ℓ with $\text{dist}(B(x_\ell^j, c_3), \partial\Omega) \notin (c_6 2^{-j}, c_7 2^{-j})$.


An oscillating u-wavelet system is called interior if it fulfils

- (further) interior support conditions, namely

$$\text{dist}(B(x_\ell^j, c_3 2^{-j}), \partial\Omega) \geq c_6 2^{-j}, j \in \mathbb{N}_0, \ell = 1, \dots, N_j.$$

Interior wavelets on Ω

The support of the first interior wavelets (on a ball):



Interior wavelet bases in $L_2(\Omega)$ - the starting point

Theorem (Triebel 2008)

Let Ω be an arbitrary domain in \mathbb{R}^n . For any $u \in \mathbb{N}_0$ there is a

$$\Phi = \left\{ \Phi_\ell^j : j \in \mathbb{N}_0, \ell = 1, \dots, N_j \right\} \subset C^u(\Omega)$$

which is

- 1 an orthonormal basis in $L_2(\Omega)$,
- 2 an interior u -wavelet system

simultaneously.

For $u = 0$ one can take the Haar Wavelet suitably restricted to Ω .

Wavelets for cubes (ii)

Let Q be the open unit cube in \mathbb{R}^n .

Theorem (Triebel 2008 - Theorem 6.30 for spaces $F_{p,q}^s(Q)$)

Let $1 < p < \infty, u, k \in \mathbb{N}_0, u > k$ and $k - \frac{m}{p} \notin \mathbb{N}_0$

for $m = 1, \dots, n$.

Then there is an oscillating u -wavelet system Φ which is a Riesz basis in the Sobolev space $W_p^k(Q)$, i.e.: An element $f \in D'(\Omega)$ belongs to $W_p^k(Q)$ if, and only if, it can be represented as

$$f = \sum_{j=0}^{\infty} \sum_{r=1}^{N_j} \lambda_r^j(f) 2^{-\frac{jn}{2}} \Phi_r^j \quad (2)$$

with λ in the sequence space $w_p^k(Q)$ and the convergence is unconditional in $W_p^k(Q)$. The representation (2) is unique with

$$\lambda_r^j(f) = 2^{jn/2} (f, \Phi_r^j).$$

Traces on the boundary of cubes (i)

Let $Q = \{x \in \mathbb{R}^n : x = (x_1, \dots, x_n), 0 < x_m < 1, m = 1, \dots, n\}$. The boundary $\Gamma = \partial Q$ of Q can be represented as

$$\Gamma = \bigcup_{\ell=0}^{n-1} \Gamma_\ell \text{ with } \Gamma_\ell \cap \Gamma_{\ell'} = \emptyset \text{ for } \ell \neq \ell',$$


where $\Gamma_\ell = \bigcup_{j=0}^{n_\ell} Q_{\ell,j}$ consists of all ℓ -dimensional faces $Q_{\ell,j}$ of Q , which are disjoint cubes of dimension ℓ .

Traces on the boundary of cubes (i)

Let $Q = \{x \in \mathbb{R}^n : x = (x_1, \dots, x_n), 0 < x_m < 1, m = 1, \dots, n\}$. The boundary $\Gamma = \partial Q$ of Q can be represented as

$$\Gamma = \bigcup_{\ell=0}^{n-1} \Gamma_\ell \text{ with } \Gamma_\ell \cap \Gamma_{\ell'} = \emptyset \text{ for } \ell \neq \ell',$$

where $\Gamma_\ell = \bigcup_{j=0}^{n_\ell} Q_{\ell,j}$ consists of all ℓ -dimensional faces $Q_{\ell,j}$ of Q , which are disjoint cubes of dimension ℓ .



Traces on the boundary of cubes (ii)

Let $tr_{\ell,j}$ be the restriction of $f \in W_p^k(\mathbb{R}^n)$ to $Q_{\ell,j}$ and

$$tr_{\ell}^r : f \mapsto TR_{\ell}^r(f) := \prod \{ tr_{\ell,j} D_{\gamma}^{\alpha} f : |\alpha| \leq r, j = 0, \dots, n_{\ell} \},$$

where only derivatives perpendicular to $Q_{\ell,j}$ are admitted.


Traces on the boundary of cubes (ii)

Let $tr_{\ell,j}$ be the restriction of $f \in W_p^k(\mathbb{R}^n)$ to $Q_{\ell,j}$ and

$$tr_{\ell}^r : f \mapsto TR_{\ell}^r(f) := \prod \{ tr_{\ell,j} D_{\gamma}^{\alpha} f : |\alpha| \leq r, j = 0, \dots, n_{\ell} \},$$

where only derivatives perpendicular to $Q_{\ell,j}$ are admitted. Then we consider the composite mapping for $\bar{r} = (r^{\ell_0}, \dots, r^{n-1})$ with $\ell_0 \leq n-1$ and $r^{\ell} = \lceil k - \frac{n-\ell}{p} \rceil - 1$:

$$tr_{\bar{r}} : f \mapsto \prod_{\ell=\ell_0}^{n-1} TR_{\ell}^{\bar{r}}(f).$$



$Q_{0,j}$: 3 directions, $\lceil k - \frac{3}{p} \rceil$ derivatives

$Q_{1,j}$: 2 directions, $\lceil k - \frac{2}{p} \rceil$ derivatives

$Q_{2,j}$: 1 direction, $\lceil k - \frac{1}{p} \rceil$ derivatives

Traces on the boundary of cubes (iii)

A u -wavelet Riesz basis cannot be interior for $k \geq 1$: Then $W_p^k(Q)$ has boundary values on the faces of Q (traces), interior wavelets do not.

Traces on the boundary of cubes (iii)

A u -wavelet Riesz basis cannot be interior for $k \geq 1$: Then $W_p^k(Q)$ has boundary values on the faces of Q (traces), interior wavelets do not. To exclude the values $k - \frac{m}{p} \notin \mathbb{N}_0$ for $m = 1, \dots, n$ is natural by the used method. The following proposition was the main part of the proof:

Proposition

Let

$$1 < p < \infty, k \neq 0 \text{ and } k - \frac{m}{p} \notin \mathbb{N}_0 \text{ for } m = 1, \dots, n.$$

Then it holds

$$\tilde{W}_p^k(Q) = \left\{ f \in W_p^k(Q) : \text{tr}_{\Gamma}^{\bar{f}} = 0 \right\},$$

Traces on the boundary of cubes (iii)

A u-wavelet Riesz basis cannot be interior for $k \geq 1$: Then $W_p^k(Q)$ has boundary values on the faces of Q (traces), interior wavelets do not. To exclude the values $k - \frac{m}{p} \notin \mathbb{N}_0$ for $m = 1, \dots, n$ is natural by the used method. The following proposition was the main part of the proof:

Proposition

Let

$$1 < p < \infty, k \neq 0 \text{ and } k - \frac{m}{p} \notin \mathbb{N}_0 \text{ for } m = 1, \dots, n.$$

Then it holds

$$\tilde{W}_p^k(Q) = \left\{ f \in W_p^k(Q) : \text{tr}_{\Gamma}^{\bar{f}} = 0 \right\},$$

Here

$$\tilde{W}_p^k(\Omega) := \{ f \in W_p^k(\mathbb{R}^n) : \text{supp } f \subset \bar{\Omega} \}.$$

The spaces on the left have **interior** u-wavelet Riesz bases.

Table of contents

- 1 Wavelets bases in function spaces on domains - the classical cases
 - Known results - wavelets on domains
 - Function spaces on domains
 - Traces on the boundary of cubes
- 2 Wavelets bases in reinforced function spaces - the exceptional values
 - The situation for cubes Q in the exceptional cases
 - Wavelet bases for reinforced function spaces on cubes Q
 - An example - $W_2^{1,\text{rinf}}(Q)$

The situation for cubes Q in the exceptional cases

Roughly speaking:

- a C^∞ -domain Ω has only boundaries of dimension $n - 1 \rightarrow$ exceptional values for $k - \frac{1}{p} \in \mathbb{N}_0$
- the cube Q has boundaries of dimension 0 to $n - 1 \rightarrow$ exceptional values for $k - \frac{m}{p} \in \mathbb{N}_0$ for $m = 1, \dots, n$

The situation for cubes Q in the exceptional cases

Roughly speaking:

- a C^∞ -domain Ω has only boundaries of dimension $n - 1 \rightarrow$ exceptional values for $k - \frac{1}{p} \in \mathbb{N}_0$
- the cube Q has boundaries of dimension 0 to $n - 1 \rightarrow$ exceptional values for $k - \frac{m}{p} \in \mathbb{N}_0$ for $m = 1, \dots, n$

Example (Grisvard '85, '92)

The space $W_2^1(Q) = F_{2,2}^1(Q)$ is exceptional: $k - \frac{2}{p} = 2 - 1$. Let $\Gamma = \partial\Omega = I_1 \cup I_2 \cup I_3 \cup I_4$. Then the trace space $tr_\Gamma W_2^1(Q)$ is the collection of all tuples $g = (g_1, g_2, g_3, g_4)$ with

$$g_\ell \in H^{\frac{1}{2}}(I_\ell), \quad \ell = 1, 2, 3, 4$$

and

$$\int_0^{1/2} \frac{|g_1(t) - g_2(t)|^2}{t} dt < \infty, \text{ etc.}$$

Reinforced function spaces for cubes Q

*Prof. Triebel: “If the mountain does not come to the prophet,
the prophet has to come to the mountain!”*

Reinforced function spaces for cubes Q

Prof. Triebel: “If the mountain does not come to the prophet, the prophet has to come to the mountain!”

Let

$$d(x) = \text{dist}(x, \partial\Omega) \text{ and } \Omega_\varepsilon := \{x \in \Omega : d(x) < \varepsilon\}.$$


Let $\Gamma_{\ell,j}$ be the ℓ -dimensional faces of the cube Q . With $\mathbb{N}_{\ell,j}^n$ we denote the multi-indices with directions perpendicular to $\Gamma_{\ell,j}$.

Definition

We say that $f \in W_p^k(\mathbb{R}^n)$ has the reinforced property $R_\ell^{r,p}$ if, and only if,

$$d^{-\frac{n-\ell}{p}} \cdot D^\alpha f \in L_p((\mathbb{R}^n \setminus \Gamma_{\ell,j})_\varepsilon) \text{ for all } \alpha \in \mathbb{N}_{\ell,j}^n, |\alpha| = r \text{ and } j = 1, \dots, n_\ell.$$

Reinforced function spaces for cubes Q (ii)



Reinforced spaces for cubes Q (iii)

Definition

Let $1 < p < \infty$, $k \neq 0$. Let

$$W_p^{k,\text{rinf}}(Q) := W_p^{k,\text{rinf}}(\mathbb{R}^n \setminus \Gamma)|_Q$$

with inf-norm and

$$W_p^{k,\text{rinf}}(\mathbb{R}^n \setminus \Gamma) :=$$

$$\left\{ f \in W_p^k(\mathbb{R}^n) : \forall 0 \leq \ell \leq n-1 : f \text{ fulfils } R_\ell^{r,p} \text{ if } r = k - \frac{n-\ell}{p} \in \mathbb{N}_0 \right\}$$

Check for every dimension ℓ if the values are exceptional and if so, add reinforce property!

Wavelet bases for reinforced function spaces on cubes Q

Theorem (main goal - for $F_{p,q}^s(Q)$ -spaces)

Let

$$1 < p < \infty, k \in \mathbb{N}_0, u > k.$$

Then there is an oscillating u -wavelet system which is a Riesz basis in $W_p^{k,\text{rinf}}(Q)$ - i. e.

$$f \in W_p^k(Q) \quad \Leftrightarrow \quad f = \sum_{j=0}^{\infty} \sum_{r=1}^{N_j} \lambda_r^j(f) 2^{-\frac{jn}{2}} \Phi_r^j$$

with $\lambda \in w_p^k(Q)$.

Wavelet bases for reinforced function spaces on cubes Q

Theorem (main goal - for $F_{p,q}^s(Q)$ -spaces)

Let

$$1 < p < \infty, k \in \mathbb{N}_0, u > k.$$

Then there is an oscillating u -wavelet system which is a Riesz basis in $W_p^{k,\text{rinf}}(Q)$ - i. e.

$$f \in W_p^k(Q) \quad \Leftrightarrow \quad f = \sum_{j=0}^{\infty} \sum_{r=1}^{N_j} \lambda_r^j(f) 2^{-\frac{jn}{2}} \Phi_r^j$$

with $\lambda \in w_p^k(Q)$.

- In the special case $k = 0$ ($L_p(\Omega)$) we can choose an interior u -wavelet system, for instance the Haar wavelet system - otherwise not.
- This theorem is a generalization of the wavelet theorem for $k - \frac{m}{p} \notin \mathbb{N}_0$ (Triebel 2008) since then there are no extra conditions.

An example - $W_2^{1,\text{rinf}}(Q) = F_{2,2}^{1,\text{rinf}}(Q)$ for $n = 2$ (i)

We have $k - \frac{2}{p} = r = 0$ - faces of dimension 0 are problematic. Hence

$$W_2^{1,\text{rinf}}(Q) = F_{2,2}^{1,\text{rinf}}(Q) = \left\{ f \in W_2^1(Q) : \int_Q |f(x)|^2 \frac{dx}{d(x)^2} \right\},$$

where d is the distance from the corner points of Q .

An example - $W_2^{1,\text{rinf}}(Q) = F_{2,2}^{1,\text{rinf}}(Q)$ for $n = 2$ (i)

We have $k - \frac{2}{p} = r = 0$ - faces of dimension 0 are problematic. Hence

$$W_2^{1,\text{rinf}}(Q) = F_{2,2}^{1,\text{rinf}}(Q) = \left\{ f \in W_2^1(Q) : \int_Q |f(x)|^2 \frac{dx}{d(x)^2} \right\},$$

where d is the distance from the corner points of Q .

Then by the theorem we find a (non-interior) oscillating u-wavelet basis which is a Riesz basis. This is not possible (in this way) for $W_2^1(Q)$ - wavelet bases (in our sense) for this space were not found yet.

The end

Thank you for your attention

Questions?