

Existence and optimal rate of convergence to equilibrium for the homogeneous Boltzmann equation

Emanuele Dolera
Università degli studi di Pavia, Italy

February 5th, 2018

Abstract

This talk deals with the spatially homogeneous Boltzmann equation for Maxwellian molecules. Conceived by L. Boltzmann in 1872 even a more general form, the equation at issue governs time evolution of the probability distribution of the particle velocities in a dilute gas. Spatial homogeneity is a suitable mathematical simplification of the complete model which highlights the effect of binary collisions on the system (rather than the transport), and the specification of “Maxwellian molecules” means we are focusing on a specific class of physical interactions between particles, whose prototype is the purely repulsive force inversely proportional to the fifth power of their distance (Maxwell force). We present three kinds of new results [1, 2]: A probabilistic representation for the solution in terms of a random sum of conditionally independent random variables, an existence result for the Boltzmann equation under the weakest assumption known so far on the collision kernel, and a definitive answer to the problem of quantifying the relaxation to equilibrium for Boltzmann-equation solutions under really mild conditions on the initial datum. This last result, which states that the total variation distance (i.e. the L^1 -norm in the absolutely continuous case) between the solution and the limiting Maxwellian distribution admits an upper bound of the form $Ce^{-\Lambda_b^* t}$, Λ_b^* being the spectral gap of the linearized collision operator and C a constant depending only on the initial datum, is at the core of kinetic theory since it provides a detailed indication about the time-scale for relaxation to equilibrium. D. Hilbert hinted at the validity of the above quantification in 1912, which was explicitly formulated as a conjecture by H.P. McKean in 1966. The main line of the new proof is based on an analogy between the problem of convergence to equilibrium and the central limit theorem of probability theory, as suggested by McKean.

References

- [1] DOLERA, E. (2015). Mathematical treatment of the homogeneous Boltzmann equation for Maxwellian molecules in the presence of singular kernels. *Ann. Mat. Pura Appl.*, **194** 1707-1732
- [2] DOLERA, E. and REGAZZINI, E. (2014). Proof of a McKean conjecture on the rate of convergence of Boltzmann-equation solutions. *Prob. Theory Relat. Fields*, **160** 315-389